



Fig. 12. Sample skeleton acquisitions of the 2-minute step test.

V. CONCLUSIONS

A multimodal dataset containing kinematic and electrophysiological data of 4 important senior fitness tests has been acquired using commercially available hardware. 11 young and 10 senior subjects have participated in the recording sessions and made their data anonymously available for research. The acquired data has sufficient quality to reproduce the scores of the different fitness tests as well as enable further studies with the available signals. Our next steps will consist in annotating specific events in the data, so temporal correlations can be assessed across different signal dimensions.

REFERENCES

- [1] Yates LBD, L.; Kurth, T.; Pong, J.E.; Gaziano T.A.M. Exceptional longevity in men: modifiable factors associated with survival and functional to age 90 years. *Arch Intern Med*. 2008;168:2550-60.
- [2] Chakrabarti S, H.B.; Ling, W.B.; Jones, J.F. Resilience, mobility and health in aging runners: a 20-year longitudinal study. *Arch Intern Med*. 2008;168:1638-46.
- [3] World Health Organization. International Classification of Functioning Disability and Health. Geneva: WHO; 2015. Available from: <http://www.who.int/classifications/icfd/en/>.
- [4] King AC. Physical activity in an aging population. *Public Health Reports*. 2008;123(2):401-26.
- [5] O'Connell K. Prevalence and correlates of physical inactivity in community-dwelling older adults in Ireland. *Prev Chronic Dis*. 2015;10(2):e0118293.
- [6] Rothman MD L-SL, Gill TM. Diagnostic significance of potentially frailty criteria. *J Am Geriatr Soc*. 2008;65(12):2211-6.
- [7] Fraix M. Role of the musculoskeletal system and the prevention of falls. *J Am Osteopath Assoc*. 2011;112(1):17-21.
- [8] Cruz-Jentoft et al. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age Ageing*. 2010 Jul;39(4):412-23. doi: 10.1093/ageing/afq034. Epub 2010 Apr 13. 2012
- [9] Ortega F. et al. Muscular strength in male adolescents and premature death: cohort study of one million participants *BMJ* 2012;345:e7279 doi: <http://dx.doi.org/10.1136/bmj.e7279>

- [10] Rikli J, Jone J (2012) Development and validation of criterion-reference clinically relevant fitness standards for maintaining physical independence in later years. *The Gerontologist*,0: 1-13. T. R., Liyanagunawardena, and S. A. Williams, S. A. "Massive open online courses on health and medicine: Review," *Journal of Medical Internet Research*, vol. 16, no. 8, e191, 2014.
- [11] Barandas M, Gamboa H, Fonseca JM, A real time biofeedback system using visual user interface for physical rehabilitation, *Procedia Manufacturing* 3, 823-828, 2015
- [12] Machado IP, Gomes AL, Gamboa H, Paixo V, Costa RM, Human activity data discovery from triaxial accelerometer sensor: Nonsupervised learning sensitivity to feature extraction parametrization, *Information Processing & Management Journal* 50, 2014
- [13] Rebelo D, Amma C, Gamboa H and Guitiz T. Human Activity Recognition for an Intelligent Knee Orthosis. In Proc. of 11th Int'l Conf. on Bio-inspired Systems and Signal Processing (BIOINSPIRAL) pp. 368-371, 2013.
- [14] Springer BA, Marin-Finckh T, Robles-Leyva G, Gill TM. Normative values for the unipedal stance test with eyes open and closed. *J Geriatr Phys Ther*, 30(1):8-15, 2007.
- [15] Jones CJ and Rose DJ (Eds), *Physical activity in later life of older adults*. Champaign, IL: Human Kinetics, 2005.
- [16] Lippincott Williams & Wilkins, *ACSM guidelines for exercise testing and prescription*, Wolters Kluwer, Ninth Edition 2014.
- [17] Roberta C. Clark and George Jones, *Senior Fitness Test Manual*, Human Kinetics (2nd Ed.), 2004.
- [18] Rose CJ, Lucchese N, Verma LD. Development of a Modified Functional Balance Scale for Use With Functionally Independent Older Adults. *Arch Phys Med Rehabil*. 2006;87(11): 1478-1485. doi:10.1016/j.apmr.2006.07.263.
- [19] Lachat E, Furrer H, Landes R and Grussenmeyer P. Assessment and calibration of the Microsoft Kinect (Kinect v2 Sensor) Towards a Potential Application for Close-Range Modeling, Remote Sensing. 2015;7:13070-13080.
- [20] Webb J and Ozkan S. Systematic Review of Kinect Applications in Elderly Age and Stroke Rehabilitation, *Journal of Neuroengineering and Rehabilitation*, 11:10.1186/s12954-014-0044-4. Disease patients," *Front. Hum. Neurosci.*, doi:10.3389/conf.fnhn.2016.220.00023, 2016.